

## The chromosomes of the rare and endemic genus *Famatinanthus* (Famatinanthoideae, Asteraceae)

## Los cromosomas del raro y endémico género *Famatinanthus* (Famatinanthoideae, Asteraceae)





## Abstract

Classical staining and *in situ* fluorescent hybridization with probes for the 18–5.8–26S and 5S genes were performed in *Famatinanthus decussatus* (endemic from Argentina), in order to know its main chromosomal characteristics and to compare them to related genera. The species is a paleopolyploid with  $2n = 54$ , likely originated from  $x=9$ , and its karyotype features are conservative: one pair of 5S signals and two of 18–5.8–26S per complement were found. All the data are congruent with the basal position of this species in the Asteraceae phylogenies.

**Keywords:** Chromosomes, endemism, *Famatinanthus decussatus*, polyploidy, rDNA.

## Resumen

Los cromosomas del raro y endémico género *Famatinanthus*. Se aplicaron tinción convencional e hibridación *in situ* fluorescente con sondas para los genes de 18–5.8–26S y 5S en *Famatinanthus decussatus* (endémica de Argentina), con el fin de conocer su características cromosómicas principales y compararlas con géneros relacionados. La especie es un paleopoliploide con  $2n = 54$ , probablemente originado a partir de  $x=9$ ; sus caracteres cariotípicos son conservados: se encontraron un par de señales de 5S por complemento y dos de 18–5.8–26S. Los datos son congruentes con la posición basal que esta especie ocupa en las filogenias de Asteraceae.

**Palabras clave:** Cromosomas, endemismo, *Famatinanthus decussatus*, poliploidía, rDNA.

## Introducción

*Famatinanthus* Ariza & S. E. Freire is a recently described genus of Asteraceae endemic to a restricted area in the province of La Rioja, Argentina (Freire *et al.*, 2014). Its single species, *F. decussatus* (Hieron.) Ariza & S. E. Freire, was formerly placed within *Aphyllocladus* Wedd. However, analyses of floral characters (i.e. corollas, anthers, style, achenes), stem anatomy, trichomes and pollen (Freire *et al.*, 2014), indicate that the species has a combination of features that do not correspond with the circumscription of *Aphyllocladus* or any other genus within the Mutisioideae tribe. Consequently, a new genus was proposed to accommodate it. The singularities of *F. decussatus* not only involve morpho-anatomical features; in fact, in a molecular phylogenetic study using 14 chloroplast DNA loci, Panero *et al.* (2014) found that the recently named genus has the two chloroplast inversions present in all Asteraceae except the nine genera of the subfamily Barnadesioideae and concluded that it is sister to the Mutisioideae–Asteroideae clade that represents more than

99% of Asteraceae. These characteristics made *Famatinanthus* to deserve a subfamily only for itself. Considering this background it is important to know the chromosomal characteristics of the species.

From the cytological point of view, chromosome counts provide a key tool in studies of systematics, phylogeny and evolution, and they are especially useful for understanding speciation and hybridization (Stebbins, 1971). Structural and quantitative characteristics of karyotypes have been significant in evolutionary and taxonomic studies in many angiosperm groups (Stebbins, 1971), being decisive in establishing linkage groups and natural classifications. (e.g. Weiss-Schneeweiss *et al.*, 2003; Pellicer *et al.*, 2010). The FISH procedure allows homologous chromosomes in a complement to be differentiated and permits the comparison among related species (Heslop-Harrison, 2000; Garcia *et al.*, 2007; Leitch *et al.*, 2008). The procedure also provides information



on genome organization and allows chromosomal evolutionary questions to be addressed (Chacón *et al.*, 2012; Pellicer *et al.*, 2010; Chiarini *et al.*, 2014). The most common molecular-cytogenetic markers are ribosomal genes (5S and 18–5.8–26S rDNA), which are abundant and highly conserved in all higher plant species (Schmidt & Heslop-Harrison, 1998). Variations in the number, signal intensity and position of rDNA loci seem to be common in several plant groups (e.g. Datson & Murray, 2006; Urdampilleta *et al.*, 2013), suggesting their mobility. Among related species, the number and location of rDNA loci may be conserved or vary considerably among populations with different ploidy levels (e.g. Adachi *et al.*, 1997; Lan *et al.*, 2011). Considering this background, the aim of our work was to describe the basic cytogenetic features of *Famatinanthus*, together with the patterns of 18–5.8–26S and 5S genes, in order to compare them to related genera.

### Materials and Methods

Seeds were bulked from plants in several natural populations in **La Rioja province**, Famatina Dpt. (S28°52'00.9" W67°41'28.1", G. Barboza *et al.* 4268; S28°50'28" W67°40'56.3", G. Barboza *et al.* 4270; S28°39'58.2" W67°42'05.5", G. Barboza *et al.* 4271). Voucher specimens are deposited at the Herbarium of the National University of Córdoba (CORD).

Mitotic chromosomes were examined in root tips obtained from seeds germinated in Petri dishes. Root tips were pre-treated in saturated p-dichlorobenzene in water for 2 h at room temperature, fixed in 3:1 ethanol/acetic acid, washed in distilled water, digested with PECTINEX® (45 min at 37 °C), and squashed in a drop of 45% acetic acid. After coverslip removal in liquid nitrogen, the slides were air dried and

stored at –20 °C. Some of these slides were used for classical staining with Giemsa. The remaining stored slides were used for determining the location and number of rDNA sites by FISH. One of the probes was the pTa71 containing the 18–5.8–26S rDNA (Gerlach & Bedbrook, 1979) labeled with biotin-14-dATP (BioNick; Invitrogen, Carlsbad, USA). For the 5S rDNA, a probe was obtained from the genome of *Prionopsis ciliata* by PCR (Moreno *et al.*, 2012) labeled with digoxigenin-11-dUTP (Roche Diagnostics). The FISH procedure was in accordance with Schwarzscher & Heslop-Harrison (2000), with minor modifications. The preparations were incubated in 100 lg/ml RNase, post-fixed in 4 % (w/v) paraformaldehyde, dehydrated in a 70–100% graded ethanol series, and air-dried. On each slide 15 µl of hybridization mixture was added (3 ng/µl of probe, 100% formamide, 50% dextran sulfate, 20 x SSC and 10% SDS), previously denatured at 70°C for 10 min. Chromosome denaturation/hybridization was done at 90°C for 10 min, 48°C for 10 min, and 38°C for 5 min using a thermal cycler (Mastercycler, Eppendorf, Hamburg, Germany), and slides were placed in a humid chamber at 37°C overnight. The probes were detected with avidin-FITC conjugate and antidigoxigenin-rodamine conjugate and counterstained and mounted with 25 µl antifade (Vectashield Vector Lab., Burlingame, USA), containing 2ng/µl of DAPI.

### Results

Seeds resulted inviable in most collected capitula. Only accession G.B. 4271 showed a good germination rate, thus the analyzed cells were all from this sample and they presented  $2n = 54$  (Fig. 1). Chromosomes are small compared to related genera of Asteraceae with  $2n = 54$ , being the average length  $c = 1.75 \pm 0.21 \mu m$  and the total



haploid length of  $LT = 94.05 \pm 2.95 \mu m$  (Table 1). The chromosomes are homogeneous in size, with a ratio between the largest and the shortest of the complement ca. 1.87. Notable secondary constrictions were detected in two chromosome pairs in most of the metaphases (Fig. 1A). Centromeres were difficult to visualize, and the high number of chromosomes hindered identifying the homologue pairs.

FISH technique with 18-5.8-26S probe evidenced four loci in terminal position, coinciding with the secondary constrictions visualized with the conventional staining, while the 5S probe showed two terminal signals per cell (Fig. 1B), located in asynteny respect to the chromosomes bearing 18-5.8-26S.

### Discussion

**Chromosome number.** As well as most basal Asteraceae, *Famatinanthus* resulted to be a high polyploid with a hypothetical basic number  $x = 9$ . Polyploids with chromosome numbers derived from  $x = 9$  have been reported for Barnadesioideae (Wulff, 1990), Mutisioideae (Ward, 1983; Waisman *et al.*, 1984; Grau, 1987) and Stifftioideae (Gibbs & Ingram, 1982) (Table 1). In fact, this number has been pointed out as basic for the entire family, which is congruent with the basal position of *Famatinanthus* within Asteraceae (Panero *et al.*, 2014). Semple & Watanabe (2009) hypothesized that multiple downward dysploid events from polyploids based on  $x = 9$  account for nearly all the base numbers reported for Mutisieae, the numbers  $x = 8$  to 32 would be the result of a long dysploid series from  $x_2 = 27$ . In this sense, *Famatinanthus* is remarkable by conserving the paleopolyploid number.


**Recurrent polyploidy.** Polyploidy is important for many aspects described

in several review articles (e.g. Soltis *et al.*, 2003; Hegarty & Hiscock, 2008; Leitch & Leitch, 2008; Van de Peer *et al.*, 2009). Autopolyploidy is a common phenomenon and it is regarded as frequent in angiosperms (Ramsey & Schemske, 1998). Within Asteraceae, Barker *et al.* (2008) revealed at least three ancient whole-genome duplications: a first one (shared by Mutisioideae, Carduoideae, Cichorioideae and Helianthoideae) placed near the origin of the family just prior to the rapid radiation of its tribes, and two independent genome duplications near the base of the tribes Mutisieae and Heliantheae. We can speculate that the genome duplication near the base of Mutisieae also affected *Famatinanthus* since they are sister clades. Thus, *Famatinanthus* would be an ancient polyploid.

**Chromosome size.** There is a phenomenon called genome downsizing whereby polyploids do not have a genome size exactly equal to the multiple of their diploid progenitors but which is somewhat smaller, as a result of a removal of redundant DNA (Leitch & Bennett, 2004). *Famatinanthus* has small chromosomes compared to hexaploid members of Barnadesioideae and Mutisioideae (Table 1). Only species of Doniophyton have  $c$  and  $LT$  values minor than *F. decussatus*. Genome size is related to the size of single chromosomes (e.g. Garnatje *et al.*, 2004), and therefore the relatively small size of the chromosomes of *F. decussatus* would be due to the longer time that the species have had from its polyploidization to rearrange its genome and eliminate redundant elements.

**rDNA sites.** Regarding Asteraceae species related to *Famatinanthus*, FISH experiments with 18-5.8-26S probe were conducted only in the Mutisioideae species





**Fig. 1.** Metaphasic chromosomes of *Famatinanthus decussatus* with (A) Giemsa staining and (B) FISH technique. Arrowheads point to NORs, arrows point to 18-5.8-26S loci (green signals) and asterisks indicate 5S loci (red signals). Bars represent 3  $\mu$ m.

*Chaptalia nutans*,  $2n = 50, 100$  (Fregonezi *et al.*, 2004) with two and four pairs of signals per complement, respectively, and also in two species of *Chaetanthera* (Baeza *et al.*, 2005a), where 1 and 3 pairs of signals in diploids with  $2n = 22$  were found. Regarding the 5S, data in species more or less related to *Famatinanthus* are known only for *Chaetanthera* (Baeza *et al.*, 2005a) in which 3 and 4 pairs of 5S signal in diploid species with  $2n = 22$  have been found. This situation evidenced the different pathways that the rDNA loci took since the two lineages split from each

other. The great variability in amount and position of rDNA loci in related species is a fact already known (Hasterok *et al.*, 2006; Heslop-Harrison & Schwarzacher, 2011; Chacón *et al.*, 2012; Morales *et al.*, 2012). For these rapid changes of copy number and chromosomal location of rDNA, different types of transposable elements have been postulated as responsible (Raskina *et al.*, 2004a, b; Belyayev *et al.*, 2005; Altinkut *et al.*, 2006; Datson & Murray, 2006).

In higher eukaryotes, the 18-5.8-26S rDNA and 5S rDNA loci are transcribed by different RNA polymerases and



usually located in different positions of chromosomes (Srivastava & Schlessinger, 1991). Colocalization of 18-5.8-26S and 5S rDNA loci has commonly been reported in animals (Dobigny *et al.*, 2003) but it is less frequent in plants (e.g. Garcia *et al.*, 2007, 2009b; Abd El-Twab & Kondo, 2006; Chang *et al.*, 2009). A dominant linked rDNA genotype was found within three large groups of Asteraceae (García *et al.*, 2010): Anthemideae, Gnaphalieae and in the “Heliantheae alliance” (Asteroideae). The remaining five tribes of the Asteroideae displayed canonical non linked arrangement of rDNA, and also the remaining 12 subfamilies, with separate organisation. *Famatinanthus* fits perfectly into this scheme, with the 5S and 18-5.8-26S loci in separate chromosomes. The results of García *et al.* (2010) indicate that nearly 25% of Asteraceae species may have developed unusual linked arrangement of rRNA genes, and the 5S gene integration within the 35S unit might have repeatedly occurred during plant evolution, and probably once in Asteraceae.

Genome downsizing, a phenomenon referred before, also affects rDNA genes. During the polyploidization process, gene reordering and gene silencing may occur (Stebbins, 1985; Soltis *et al.*, 2003; Leitch & Bennett, 2004; Pires *et al.*, 2004). In Solanaceae, for example, an analysis of the 18-5.8-26S rDNA of *Nicotiana* indicated that parental loci were initially maintained in newly formed polyploids, although the sequences within a locus might be subject to concerted evolution, and over periods greater than 1 myr, individual loci would disappear (Kovářík *et al.*, 2008). In Asteraceae, a similar situation was found in *Tragopogon* (Buggs *et al.*, 2009; Chester *et al.*, 2012).

Given that *Famatinanthus* is a

hexaploid, more than a pair of 5S signals per complement could be expected. Instead, only a pair was found, which would be an indication of an ancient polyploidy. It has been found that the number of loci of 5S is subject to dynamic, rapid, changes (cfr. sup.). Loss of gene copies might be taking place at these loci, as demonstrated in members of other several plant families (i.e. Kotseruba *et al.* 2003; Renny-Byfield *et al.*, 2012). A similar situation was observed in polyploid species in other Asteraceae genera such as *Brachyscome* (Asteroideae, Adachi *et al.*, 1997), *Artemisia* (Asteroideae, Garcia *et al.*, 2009a; Pellicer *et al.*, 2013), *Xeranthemum* (Carduoideae, Garnatje *et al.*, 2004), *Tragopogon* (Cichorioideae, Buggs *et al.*, 2009; Chester *et al.*, 2012), in which the number of 18-5.8-26S sites seemed to evolve faster than the 5S sites, as they do not increase with the successive genome additions. By contrast, genome sizes of some polyploid taxa present additivity, although data pointing to genome upsizing in older polyploids are limited (Garcia *et al.*, 2009; Leitch *et al.*, 2008).

FISH studies with rDNA probes would be desirable in genera more closely related to *Famatinanthus* (e.g. *Barnadesia*, *Duseniella*, *Dasyphyllum*) in order to achieve a more complete picture of chromosomal evolution in basal Asteraceae.

### Acknowledgements

The authors thank Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina), SECyT-UNC, and SECyT-UNRC for financial support. Thanks are also tendered to

D. Olivera for his unvaluable assistance in Famatina's field work (La Rioja).



# Literature cited

- Abd El-Twab, M. H. & K. Kondo.** 2006. FISH physical mapping of 5S, 45S and Arabidopsis-type telomere sequence repeats in *Chrysanthemum zawadskii* showing intra-chromosomal variation and complexity in nature. *Chromosome Bot.*, 1: 1-5.
- Adachi, J.; K. Watanabe; K. Fukui; N. Ohmido & K. Kosuge.** 1997. Chromosomal location and reorganization of the 45S and 5S rDNA in the *Brachyscome lineariloba* complex (Asteraceae). *J. Plant Res.*, 110: 371-377.
- Altinkut, A.; V. Kotseruba; V. M. Kirzhner; E. Nevo; O. Raskina & A. Belyayev.** 2006. Ac-like transposons in populations of wild diploid Triticeae species: comparative analysis of chromosomal distribution. *Chromosome Res.*, 14: 307-317.
- Baeza, C.; E. Ruiz & M. Negritto.** 2008. The karyotype of *Chaetanthera incana* Poepp. (Asteraceae). *Gayana Bot.*, 65: 237-240.
- Baeza, C. & C. Torres-Díaz.** 2006. The karyotype of *Chaetanthera pentacaenoides* (Phil.) Hauman (Asteraceae). *Gayana Bot.*, 63: 180-182.
- Baeza, C. M. & O. Schrader.** 2005a. Análisis del cariotipo y detección de los genes 5S y 18S/25S rDNA en *Chaetanthera microphylla* (Cass.) Hook. et Arn. (Asteraceae). *Gayana Bot.*, 61: 49-51.
- Baeza, C. & O. Schrader.** 2005b. Karyotype analysis in *Chaetanthera chilensis* (Willd.) DC. and *Chaetanthera ciliata* Ruiz et Pavón (Asteraceae) by double fluorescence in situ hybridization. *Caryologia*, 58: 332-338.
- Baeza, C. M.; E. Ruiz; P. Novoa & M. A. Negritto.** 2010a. El cariotipo de *Chaetanthera linearis* Poepp. (Asteraceae). *Gayana Bot.*, 67: 113-116.
- Baeza, C. M.; E. Ruiz & C. Torres-Díaz.** 2010b. El cariotipo de *Chaetanthera renifolia* (J. Remy) Cabrera (Asteraceae). *Gayana Bot.* 67: 246-248.
- Barker, M. S.; N. C. Kane; M. Matvienko; A. Kozik; R. W. Michelmore; S. J. Knapp & L. H. Rieseberg.** 2008. Multiple paleopolyploidizations during the evolution of the Compositae reveal parallel patterns of duplicate gene retention after millions of years. *Mol. Biol. Evol.*, 25: 2445-2455.
- Buggs, R. J. A.; A. N. Doust; J. A. Tate; J. Koh; K. Soltis; F. A. Feltus; A. H. Paterson; P. S. Soltis & D. E. Soltis.** 2009. Gene loss and silencing in *Tragopogon miscellus* (Asteraceae): comparison of natural and synthetic allotetraploids. *Heredity*, 103: 73-81.
- Chacón, J.; A. Sousa; C. M. Baeza & S. S. Renner.** 2012. Ribosomal DNA distribution and a genus-wide phylogeny reveal patterns of chromosomal evolution in K (Alstroemeriaceae). *Am. J. Bot.*, 99: 1501-1512.
- Chang, Y. C.; C. T. Shii & M. C. Chung.** 2009. Variations in ribosomal RNA gene loci in spider lily (*Lycoris* spp.). *J. Am. Soc. Horti. Sci.*, 134: 567-573.
- Chiarini, F. E.; F. F. Santiñaque; J. D. Urdampilleta & M. L. Las Peñas.** 2014. Genome size and karyotype diversity in *Solanum* sect. Acanthophora (Solanaceae). *Plant Syst. Evol.*, 300(1): 113-125.
- Cialdella, A. M. & A. G. López de Kiesling.** 1981. Cariología de *Schlechtendahlia luzulaefolia* Less. (Compositae). *Darwiniana*, 23: 357-360.
- Datson, P. M. & B. G. Murray.** 2006. Ribosomal DNA locus evolution in *Nemesia*: transposition rather than structural rearrangement as the key mechanism? *Chromosome Res.*, 14: 845-857.
- Dobigny, G.; C. Ozouf-Costaz; C. Bonillo & V. Volobouev.** 2003. Evolution of rRNA gene clusters and telomeric repeats during explosive genome repatterning in *Taterillus* X (Rodentia, Gerbillinae). *Cytogenet. Genome Res.*, 103(1-2): 1-10.



- 2): 94-103.
- Fregonezi, J. N.; J. Torezan & A. L. Vanzela.** 2004. A karyotypic study of three southern Brazilian Asteraceae species using fluorescence in situ hybridization with a 45S rDNA probe and C-CMA3 banding. *Genet. Mol. Biol.*, 27: 223-227.
- Freire, S. E.; G. E. Barboza; J. J. Cantero & L. Ariza Espinar.** 2014. *Famatinanthus*, a new Andean genus segregated from *Aphyllocladus* (Asteraceae). *Syst. Bot.*, 39: 349-360.
- Fukushima, K.; K. Imamura; K. Nagano & Y. Hoshi.** 2011. Contrasting patterns of the 5S and 45S rDNA evolutions in the *Byblis liniflora complex* (Byblidaceae). *J. Pl. Res.*, 124: 231-244.
- Garcia, S.; T. Garnatje; O. Hidalgo; E. D. Mcarthur; S. Šiljak-Yakovlev & J. Vallès.** 2007. Extensive ribosomal DNA (18S–5.8S–26S and 5S) colocalization in the North American endemic sagebrushes (subgenus *Tridentatae*, *Artemisia*, Asteraceae) revealed by FISH. *Pl. Syst. Evol.*, 267: 79–92.
- Garcia, S.; T. Garnatje; J. Pellicer; E.D. Mcarthur; S. Šiljak-Yakovlev & J. Vallès.** 2009a. Ribosomal DNA, heterochromatin, and correlation with genome size in diploid and polyploid North American endemic sagebrushes (*Artemisia*, Asteraceae). *Genome*, 52: 1012–1024.
- Garcia, S.; K. Y. Lim; M. Chester; T. Garnatje; J. Pellicer; J. Vallès & A. Kovařík.** 2009b. Linkage of 35S and 5S rRNA genes in *Artemisia* (family Asteraceae): first evidence from angiosperms. *Chromosoma*, 118: 85-97.
- Garcia, S.; J. L. Panero; J. Siroky & A. Kovařík.** 2010. Repeated reunions and splits feature the highly dynamic evolution of 5S and 35S ribosomal RNA genes (rDNA) in the Asteraceae family. *BMC Pl. Biol.*, 10: 176.
- Garnatje, T.; J. Vallès; S. Garcia; O. Hidalgo; M. Sanz; M. A. Canela & S. Šiljak-Yakovlev.** 2004. Genome size in *Echinops* L. and related genera (Asteraceae, Cardueae): karyological, ecological and phylogenetic implications. *Biol. Cell*, 96:117–124
- Gerlach, W. L. & J. R. Bedbrook.** 1979. Cloning and characterization of ribosomal RNA genes from wheat and barley. *Nucleic Acids Res.*, 7: 1869-1885.
- Gibbs, P. E. & R. Ingram.** 1982. Chromosome numbers of some Brazilian flowering plants. *Notes Roy. Bot. Gard. Edinburgh*, 40: 399–407.
- Grau, J.** 1987. Chromosomenzahlen chilenischer Mutisieen (Compositae). *Bot. Jahrb. Syst.*, 108: 229–237.
- Grubbs, K. C.; R. L. Small & E. E. Schilling.** 2009. Evidence for multiple, autopoloid origins of agamosperous populations in *Eupatorium sessilifolium* (Asteraceae). *Pl. Syst. Evol.*, 279: 151-161.
- Hegarty, M. J. & S. J. Hiscock.** 2008. Genomic clues to the evolutionary success of polyploid plants. *Curr. Biol.*, 18: 435–444.
- Heslop-Harrison, J. S. & T. Schwarzacher.** 2011. Organisation of the plant genome in chromosomes. *Plant J.*, doi:10.1111/j.1365-313X.2011.04544.x
- Heslop-Harrison, J. S.** 2000. Comparative genome organization in plants: from sequence and markers to chromatin and chromosomes. *Pl. Cell*, 12: 617-635.
- Kotseruba, V.; D. Gernand; A. Meister & A. Houben.** 2003. Uniparental loss of ribosomal DNA in the allotetraploid grass *Zingieria trichopoda* (2n = 8). *Genome*, 46: 156-163.
- Kovařík, A.; M. Dadejova; Y. K. Lim; M. W. Chase; J. J. Clarkson; S. Knapp & A. R. Leitch.** 2008. Evolution of rDNA in *Nicotiana* allopolyploids: a potential link between rDNA homogenization and epigenetics. *Ann. Bot.*, 101: 815-823.
- Lan, T. & V. A. Albert.** 2011. Dynamic distribution patterns of ribosomal DNA and chromosomal evolution in *Paphiopedilum*, a lady's slipper or-




- chid. BMC Plant Biol., 11: 126.
- Leitch, A. R. & I. J. Leitch.** 2008. Genomic plasticity and the diversity of polyploid plants. *Science*, 320: 481–483.
- Leitch, I. J.; L. Hanson; K. Y. Lim; A. Kovařík; M. W. Chase; J. J. Clarkson, & A. R. Leitch.** 2008. The ups and downs of genome size evolution in polyploid species of *Nicotiana* (Solanaceae). *Ann. Bot.*, 101(6): 805–814.
- Morales A. G.; M. L. R. Aguiar-Perecin & M. Mondin.** 2011. Karyotype characterization reveals an up and down of 18-5.8-26S and 5S rDNA sites in *Crotalaria* (Leguminosae-Papilionoideae) species of the section Hedriocarpae subsection Macrostachyae. *Gen. Res. Crop. Evol.* DOI 10.1007/s10722-011-9683-8
- Moreno, N. C.; L. Stiefkens; M. L. Las Peñas; A. Bartoli; R. Tortosa & G. Bernardello.** 2012. Molecular cytogenetic studies of the “Xanthocephalum group” (Asteraceae). *Pl. Syst. Evol.*, 298: 1503–1514.
- Panero, J. L.; S. E. Freire; L. Ariza Espinar; B. S. Crozier; G. E. Barboza & J. J. Cantero.** 2014. Resolution of deep nodes yields an improved backbone phylogeny and a new basal lineage to study early evolution of Asteraceae. *Mol. Phylogenet. Evol.*, 80: 43–53.
- Pellicer, J.; T. Garnatje; J. Molero; F. Pustahija; S. Šiljak-Yakovlev & J. Vallès.** 2010. Origin and evolution of the South American endemic *Artemisia* species (Asteraceae): evidence from molecular phylogeny, ribosomal DNA and genome size data. *Aust. J. Bot.*, 58: 605–616.
- Pellicer, J.; S. Garcia; J. Vallès; K. Kondo & T. Garnatje.** 2013. FISH mapping of 35S and 5S rRNA genes in *Artemisia* subgenus *Dracunculus* (Asteraceae): changes in number of loci during polyploid evolution and their systematic implications. *Bot. J. Linnean Soc.*, 171: 655–666.
- Pires, J.; C., J. Zhao; M. Schranz; E. J. Leon; P. A. Quijada; L. N. Lukens & T. C. Osborn.** 2004. Flowering time divergence and genomic rearrangements in resynthesized *Brassica* polyploids (Brassicaceae). *Biol. J. Linnean Soc.*, 82: 675–688.
- Ramsey, J. & D. W. Schemske.** 1998. Pathways, mechanisms and rates of polyploid formation in flowering plants. *Annu. Rev. Ecol. Syst.*, 29: 467–501.
- Raskina, O.; A. Belyayev & E. Nevo.** 2004a. Activity of the En/Spm-like transposons in meiosis as a base for chromosome repatterning in a small, isolated, peripheral population of *Aegilops speltoides* Tausch. *Chromosome Res.*, 12: 153–161.
- Raskina, O.; A. Belyayev & E. Nevo.** 2004b. Quantum speciation in *Aegilops*: molecular cytogenetic evidence from rDNA cluster variability in natural populations. *Proc. Natl. Acad. Sci. U.S.A.*, 101: 14818–14823.
- Renny-Byfield, S.; A. Kovařík; M. Chester; R. A. Nichols; J. Macas; P. Novák & A. R. Leitch.** 2012. Independent, rapid and targeted loss of highly repetitive DNA in natural and synthetic allopolyploids of *Nicotiana tabacum*. *PloS One*, 7: e36963.
- Schmidt, T. & J. S. Heslop-Harrison.** 1998. Genomes, genes and junk: the large-scale organization of plant chromosomes. *Trends Pl. Sci.*, 3: 195–199.
- Schwarzacher, T. & P. Heslop-Harrison.** 2000. Practical in situ hybridization. Bios Scientific Publishers Limited, Oxford.
- Semple, J. C. & K. Watanabe.** 2009. A review of chromosome numbers in Asteraceae with hypotheses on chromosomal base number evolution. In: Funk, V. A.; A. Susanna; T. F. Stuessy & R. J. Bayer (eds.) *Systematics, evolution, and biography of Compositae*. IAPT, Vienna, pp. 61–72.
- Soltis, D. E.; P. S. Soltis, M. D. Bennett & I. J.**



- Leitch.** 2003. Evolution of genome size in the Angiosperms. *Am. J. Bot.*, 90: 1596–1603.
- Srivastava, A. K., & D. Schlessinger.** 1991. Structure and organization of ribosomal DNA. *Biochimie*, 73(6): 631-638.
- Stebbins, G. L.** 1971. Chromosomal evolution in higher plants. E. Arnold, London.
- Stebbins, G. L.** 1985. Polyploidy, hybridization, and the invasion of new habitats. *Ann. Missouri Bot. Gard.*, 72: 824-832.
- Van de Peer, Y.; S. Maer & A. Meyer.** 2009. The evolutionary significance of ancient genome duplications. *Nat. Rev. Genet.*, 10: 725–732.
- Waisman, C. E.; E. Rozenblum & J. H. Hunziker.** 1984. Estudios cariológicos en Compositae. I. *Darwiniana*, 25: 217-226.
- Ward, D. E.** 1983. Chromosome counts from New Mexico and southern Colorado. *Phytologia*, 54(5): 302-308.
- Watanabe, K.; T. Yahara; G. Hashimoto; Y. Nagatani; A. Soejima; T. Kawahara & M. Nakazawa.** 2007. Chromosome numbers and karyotypes in Asteraceae 1. *Ann. Missouri Bot. Gard.*, 94(3): 643-654.
- Weiss-Schneeweiss, H.; T. F. Stuessy; S. Šiljak-Yakovlev; C. M. Baeza & J. Parker.** 2003. Karyotype evolution in South American species of *Hypochaeris* (Asteraceae, Lactuceae). *Pl. Syst. Evol.*, 241(3-4): 171-184.
- Wulff, A. F.** 1990. Chromosome studies in Barnadesiinae (Mutisieae, Asteraceae). I. *Chuquiraga* and *Doniophyton*. *Darwiniana*, 30 (1-4): 185.
- Wulff, A. F.** 1995. El cariotipo de *Barnadesia odorata* (Asteraceae). *Hickenia*, 2(24-31): 129-130.





**Fig. 2.** *Famatinanthus decussatus*. A. Hábitat; B. Planta mostrando las raíces; C. Tallo; D. Capítulo; E. Capitulecencia; F. Cipselas. (Fotografías G. Barboza, J. Cantero; R. Deanna & S. Leiva 4270, CORD, HAO).



